


# ● CHROMagar™ Staph aureus



For isolation and direct differentiation  
of *Staphylococcus aureus*

**CHR**  **Magar**  
The Chromogenic Media Pioneer



## For isolation and direct differentiation of *Staphylococcus aureus* in clinical and industrial samples

### Background

**Food Industry:** Human beings are the main reservoir of *S.aureus*. A carrier contaminates the surrounding environment when coughing, sneezing and by touching food with a hand having a *staphylococcus*-infected lesion. It is often found in the environment and on food preparation surfaces and also in certain uncooked foods (dairy products, salads, sandwiches...). It is important to check the presence of *S.aureus* before and after the foodstuff sterilisation process.

**Clinical relevance:** *S.aureus* is the leading cause of skin and soft tissue infections and can also cause serious infections such as bloodstream infections, pneumonia, or bone and joint infections.

### Medium Performance

#### Clinical application

- EASY TO READ**  
compared to Blood Agar or Mannitol Salt Agar. CHROMagar™ Staph aureus allows easier differentiation of *S. aureus* colonies enhanced by a mauve colour and is of considerable help in identifying suspect colonies. Thus, it reduces the confirmatory workload.

- HIGH SENSITIVITY**

exceeds 99%\*

\*Specificity and sensibility from scientific study: "Evaluation of CHROMagar Staph aureus, a new chromogenic medium, for isolation and presumptive identification of *Staphylococcus aureus* from human clinical specimens." Gaillot O. et al. 2001. Journal of Clinical Microbiology, 38: 1587-1591.

\*\*\*

#### Food and environmental quality control


- EASY TO PREPARE**  
The conventional medium for *S.aureus* is the Baird-Parker which has to be supplemented with RPF (Rabbit Plasma Fibrinogen), rendering the plate manufacturing delicate and complex, and also reducing the shelf life of the poured plates to a couple of weeks. On the contrary, CHROMagar™ Staph aureus comes with all the compounds already in the agar (no need of any supplement).
- FAST**  
The results on Baird Parker have to be read after 48h of incubation while with CHROMagar™ Staph aureus the results are available after only 24h.

### Medium Description

<b>Powder Base</b>	Total .....	82.5 g/L
	Agar .....	15.0
	Peptone and yeast extract .....	40.0
	Salts .....	25.0
	Chromogenic mix .....	2.5
	Storage at 15/30°C - pH: 6.9 +/- 0.2	
	Shelf Life .....	2 years

Usual Samples	Clinical: wounds, sputum Industrial: Food stuff
Procedure	Direct streaking. Incubate at 37°C for 18-24 h. Aerobic conditions.

Scientific Publications on this product: available on [www.CHROMagar.com](http://www.CHROMagar.com)  
For detailed preparation procedure, please refer to our IFU.



#### Plate Reading

- *S. aureus*  
→ pink to mauve
- Other bacteria  
→ colourless, blue or inhibited



### Quality Control Strains

<i>S. aureus</i> ATCC® 25923 .....	mauve
<i>S. saprophyticus</i> ATCC® 15305 .....	turquoise blue
<i>E. coli</i> ATCC® 25922 .....	inhibited
<i>C. albicans</i> ATCC® 60193 .....	inhibited
<i>E. faecalis</i> ATCC® 29212 .....	inhibited

ATCC® is a registered trademark of the American Type Culture Collection

### Order References

Please use these product references when contacting your local distributor:

- 1000 ml pack ..... TA670
- 5000 ml pack ..... TA672
- Bulk ..... on request

CHROMagar  
4 place du 18 juin 1940  
75006 Paris - France

Find your nearest distributor on  
[www.CHROMagar.com/contact](http://www.CHROMagar.com/contact)